DEVELOPMENT OF TWO MOBILE MEASUREMENT STATIONS FOR AMBIENT AIR TOXIC MONITORING IN COMMUNITIES

Andrea Polidori*



South Coast Air Quality Management District (AQMD) 21865 Copley Dr, Diamond Bar, CA 91765

INTRODUCTION

Background

• Multi-year Community-Scale Air Toxics Monitoring grant from EPA (November 2008)


Specific Objectives

- Develop two rapidly deployable mobile measurements stations (MMS) for ambient air toxics monitoring
- Characterize ambient air toxics levels in communities surrounding LAX and LGB
- Supplement a concurrent LAWA air quality study by providing additional capabilities for air toxics monitoring

MMS Characteristics


- Fast-response deployment
 - Wide spatial coverage and rapid adjustment
- Flexibility of operation
 - Land-based power and self-contained generator or battery power
- Gaseous and particulate pollutants characterization
- Multiple monitoring technologies
 - Continuous (e.g. 1- to 5-min)
 - Integrated (e.g. 24-hr)
- Address AQMD and community concerns
 - Local impacts of pollution sources

Particulate Matter \



Air flow (adjustable)


Air conditioner



HVAC duct , (8" x 16")











Six port manifold

Power options: Shore power, Battery bank, and Propane generator; (Solar panels)

Instrumentation

- Continuous
 - DustTrak DRX (TSI, Inc.): PM₁, PM_{2.5}, PM₄, PM₁₀, and TPM (μg/m³)
 - w-CPC (TSI, Inc.): UFP (#/cm³)
 - Aethalometer (Magee, Inc.): BC (μg/m³)
 - NOx Monitor (2B Technologies): NO, NO₂, and NOx (ppb)
 - Baseline-Mocon Series 9000 (Mocon, Inc.): Methane and NMHC (ppm)
 - Syntech Spectras GC (Syntech, Inc.): Individual organic compounds
 - Q-Trak (TSI, Inc.): CO and CO₂
 - Weather Station (Climatronics, Inc.): T, RH, WS, WD, and P
- Integrated
 - BGI (Omni, Inc.) filter-based PM sampler
 - Speciated analysis
 - ◆ Xontech multi-canister sampling (Xontech, Inc.)
 - VOCs
















AIT TOXICS MONITORING IN COLTON





- Sampling location: proposed recreational park
- Sampling period: 02/16/10 to 04/07/10
- Objective: determine I-10 / I-215 emission contribution s
- Pollutants measured: PM₁₀, UFP, and BC




AIT TOXICS MONITORING IN COLTON





- Sampling location: proposed recreational park
- Sampling period: 02/16/10 to 04/07/10
- Objective: determine I-10 / I-215 emission contribution s
- Pollutants measured: PM₁₀, UFP, and BC






AIT TOXICS MONITORING IN COLTON: RESULTS






- PM_{10} Colton $< PM_{10}$ San Bernardino $< PM_{10}$ Fontana (background) $< PM_{10}$ MATES III (37.0 $\mu g/m^3$) <<< NAAQS (150 $\mu g/m^3$)
- UFP Colton < UFP San Bernardino < UFP Fontana (background) < UFP South Coast Air Basin


AIT TOXICS MONITORING IN COLTON: RESULTS



- BC Colton ~ BC South Coast Air Basin
- BC at all three sampling sites tracked each other well



FUTURE WORK: LGB Airport Study Plan



Objectives

- Air toxics levels in LGB communities
- Concentration gradients driven by proximity to LGB
- Area impacted by aircraft emissions and airport activities
- Freeway contributions
- Baseline data for longer term measurements

Part of EPA funded community-scale air monitoring grant

SUMMARY

- Two MMS for fast response deployment in communities of the South Coast Air Basin were successfully developed
- One of the MMS was tested at a proposed recreational park in Colton. Measured air pollution levels were typical for the region
- A follow-up study will be conducted in 2011 during different seasonal conditions
- The two MMS will soon be deployed at LGB and LAX

ACKNOWLEDGEMENTS

Steve Boddeker, Richard Parent and the SCAQMD Special Monitoring group